

INVERT CONTAINER AND SHAKE WELL BEFORE USE

KEEP OUT OF REACH OF CHILDREN

INVERT CONTAINER AND SHAKE WELL BEFORE USE

biostart®



HSNO Classifications: 6.4A, 6.9B, 9.1D

**Warning:** Terracin is mildly irritating to the skin and irritating to the eye. When handling Terracin wear protective clothing, glasses and gloves and avoid contact with skin and eyes. Terracin is a soil biocide and is toxic to the aquatic environment, so avoid product release to the environment. Dispose of used container carefully.

**DIRECTIONS FOR USE. READ LABEL BEFORE USE.**

**Mixing Instructions:** Partially fill the spray tank with clean water and start agitation. Add the required amount of Terracin to the tank. Complete filling the tank to the required volume. **Handling precautions:** When mixing or applying, avoid contact with skin and eyes. Wear protective clothing, gloves and goggles. Flush spray equipment thoroughly with water after use. Do not eat or drink while using. Wash protective clothing after use. **Application:** For best results apply to damp soil or water in. Sprayers should be calibrated before use. Wash hands and face thoroughly after handling. Avoid release to the environment. Do not apply onto or into waterways. **Compatibility:** Do not mix Terracin with Mycorrcin or Digester. **Storage instructions:** Store below 30°C. Keep out of reach of children. Keep container tightly closed. Do not store diluted product. **Empty container disposal:** Triple rinse and return to an approved recycling centre. **First Aid:** If medical advice is needed have the product container or label on hand. If skin contact occurs, remove contaminated clothing and flush skin with running water, if irritation occurs seek medical advice. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists seek medical advice. **Spills:** When dealing with spills wear personal protective clothing and equipment. Contain the spill by damming, recover spilt product then transfer the recovered spilt material to a properly labelled drum. Deal with all spillages immediately to avoid product being released to the environment.

For more information call: **New Zealand: 0800 116 229, Australia: 1800 359 559**

biostart®



Manufactured & distributed in New Zealand by: BioStart Limited. Ph 0800 116 229 [www.biostart.co.nz](http://www.biostart.co.nz) [info@biostart.co.nz](mailto:info@biostart.co.nz)

Australian distributor: BioStart Brands Pty Ltd. Ph 1800 359 559 [www.biostart.com.au](http://www.biostart.com.au)



# Terracin™

## SOIL BIOCIDES

Contains BS 1b which has a broad range of anti-fungal activity

Active: *Bacillus Amyloliquefaciens* BS 1b not less than 1 x 10<sup>9</sup>cfu/L



Store out of direct sunlight and below 30°C

net contents **20 L**

Made in New Zealand

CROP	TIMING	TYPE OF APPLICATION & RATE	RECOMMENDED FOLLOW-UP PROGRAMME
Kiwifruit	Early winter	<b>Orchard floor:</b> Early winter apply 20 L/ha Terracin, water rate 1,000 L/ha, (1:50 dilution) to the orchard floor.	<b>Orchard floor:</b> 3 weeks after Terracin application apply 10 L/ha Digester in 1,000 L water. In early spring apply 10 L/ha Mycorrcin in 1,000 L water.
	For orchards prone to being wet over winter	<b>Spot spray treatment for infected vines:</b> Early winter apply to the root zone 20L/vine a solution of 20L Terracin/1,000L water (1:50 dilution). Treat all adjacent vines at the same rate.	<b>Spot spray treatment for infected vines:</b> 3 weeks after Terracin application apply to the root zone 20 L/vine a solution of 10 L Digester/1,000 L water (1:100 dilution). Treat all adjacent vines at the same rate. In early summer and at autumn root flush apply 5 L/ha Mycorrcin in 1,000 L water.
	Post sluicing:	Following sluicing apply 20 L/hole a solution of 20 L Terracin/1,000L water (1:50 dilution).	<b>Post sluicing:</b> 3 weeks after Terracin application apply 20L/hole a solution of 10 L Digester/1,000 L water (1:100 dilution). In early summer and at autumn root flush apply 5 L/ha Mycorrcin in 1,000 L water.
Pipfruit, Stonefruit, Citrus, Grapes, Berryfruit and Vegetables	Late winter/early spring	<b>Orchard floor:</b> Late winter/early spring apply 20 L/ha Terracin, water rate 1,000L/ha, (1:50 dilution) to the orchard floor.	<b>Orchard floor:</b> 3 weeks after Terracin application apply 10 L/ha Mycorrcin in 1,000 L water. In early summer and at autumn root flush apply 5 L/ha Mycorrcin in 1,000 L water.
	For orchards in drier areas	<b>Spot spray treatment for infected vines:</b> Late winter/early spring apply to the root zone 20 L/vine a solution of 20L Terracin/1,000 L water (1:50 dilution). Treat all adjacent vines at the same rate.	<b>Spot spray treatment for infected vines:</b> 3 weeks after Terracin application apply to the root zone 20 L/vine a solution of 10 L Mycorrcin/1,000 L water (1:100 dilution). Treat all adjacent vines at the same rate. In early summer and at autumn root flush apply 5 L/ha Mycorrcin in 1,000 L water.
	Post sluicing:	Following sluicing apply 20 L/hole a solution of 20 L Terracin/1,000 L water (1:50 dilution).	<b>Post sluicing:</b> 3 weeks after Terracin application apply 20 L/hole a solution of 10 L Mycorrcin/1,000 L water (1:100 dilution). In early summer and at autumn root flush apply 5 L/ha Mycorrcin in 1,000 L water.